

Is there a case to allow snares to be used to undertake scientific research on foxes in Scotland?

Professor Stephen Harris BSc PhD DSc

Background

The National Anti Snaring Campaign asked me to assess whether there is a case to permit the use snares to catch foxes for research purposes in Scotland.

In considering this issue, I have reviewed how often snares have been used to catch foxes for research in Scotland, the animal welfare implications of using snares to catch foxes for research, the ethical implications of using snares for research, and whether any study that used snares to catch foxes for research would be likely to meet the ethical standards for publication in a reputable scientific journal.

How often have snares been used for scientific research in Scotland?

To assess the need to use snares to undertake research on foxes in Scotland, I considered how often they have been used to catch foxes for research hitherto. Since fox snares only came into use in the 1960s, following the development of small-diameter steel cables (Harris 2022), I have only considered research published from 1970 onwards.

There is a long history of scientific research into all aspects of fox behaviour, biology and ecology in Scotland. I have listed the first hundred papers published since 1970 that came up in a search on Google Scholar (see Appendix). I have only included peer-reviewed primary studies, i.e., review papers were excluded. While most of the studies listed in the Appendix were based primarily or exclusively in Scotland, I have included studies where data collected in Scotland made a significant contribution to a wider British or UK study. While not comprehensive, this list of peer-reviewed research provides an overview of the types of studies undertaken on foxes in Scotland over the past 53 years, and how often snares were used to catch foxes for research.

Studies undertaken in the 1970s and 1980s showed that the major influence on fox population dynamics was food supply, not anthropogenic population control. More recent studies have included studies on the number and distribution of foxes in Scotland, the economic impact of fox predation on sheep, parasites of economic importance, and the impact of fox predation on ground-nesting birds. One study, not included in the Appendix because it was not on foxes *per se*, looked at the negative impacts of snaring foxes on capercaillie numbers (Grove & Oswald 2001).

None of these studies used snares to catch foxes for their research. Many of the studies into the impacts of predator control on ground-nesting bird populations relied on existing predator control measures that were already being undertaken undertaken in the area but the papers were unclear about the techniques that were

used, or the relative contribution of different predator control methods. For instance, in one study *fox, stoat, weasel, and some corvid species ... were routinely killed year-round by full-time gamekeepers employed to maximise numbers of red grouse. Foxes were either shot by a rifle at night or caught and restrained by metal snares* (Appendix: reference 7). In one case, where predator control was undertaken specifically for the study, professional contractors located foxes using infra-red or thermal imaging devices and shot them using a rifle. Snares were not used (Appendix: reference 16).

So there is no need to use snares for studies of the impacts of predators on ground-nesting birds. In fact, other techniques that are available are more acceptable to practitioners, who rated shooting at night with a rifle significantly more effective than other methods of killing foxes. Practitioners considered snares to be the least acceptable way of catching and killing foxes (White *et al.* 2003).

Over the last half-century, a wide variety of research techniques have been employed to study foxes in Scotland, and a number of new research techniques, such as molecular genetics, stable isotopes, video recording and camera trapping, and identifying individual foxes from their physical appearance (Dorning & Harris 2019), have become available to scientists. None of these techniques require that a fox is caught.

Despite the number, and diversity, of studies undertaken on foxes in Scotland over the last 53 years, snares have rarely, if ever, been used to catch foxes for research purposes. There is no evidence that snares will be needed for future research studies, and a variety of new research techniques are becoming available to wildlife biologists that do not involve catching foxes.

Welfare issues associated with using snares for research

The principal law relating to animal welfare in Scotland (and the rest of the UK) is the Animal Welfare Act 2006 (<https://www.gov.uk/guidance/animal-welfare#legislation>). Among other things, this Act prohibits causing unnecessary suffering to a captive animal. Once an animal is caught in a snare, irrespective of whether it is a target or non-target species, it is captive and is covered by this Act.

The British Veterinary Association and the British Veterinary Zoological Society (a specialist group of the British Veterinary Association) issued a position statement in which they state that *Due to the nature of snares and the duration of time animals may legally be held in snares even when best practice is followed, the potential negative animal welfare impacts are significant.* They went on to say that *The speed at which welfare begins to be impacted is rapid (seconds from the moment of restraint) and suffering can be prolonged, that in some cases [this occurs] over a considerable length of time, and that snares can also be indiscriminate and may result in the capture and suffering of non-target species.* In view of their very significant welfare concerns over the use of snares, the British Veterinary Association and the British Veterinary Zoological Society called on the UK

governments to introduce *An outright ban on the use and sale of snares to both the general public and trained operators* (British Veterinary Association 2022, <https://www.bva.co.uk/take-action/our-policies/snaring/>).

The role of the Scottish Animal Welfare Commission (SAWC) is to provide advice on animal welfare, including, specifically, consideration of how policies take account of animal sentience, the wider welfare needs of animals and the type of improvements that could be made. They also concluded *that snares cause significant welfare harms to members of both target and non-target species*, and recommended that *the sale of snares and their use by both public and industry are banned in Scotland, on animal welfare grounds* (Scottish Animal Welfare Commission 2022).

In this respect, it should be remembered that, while many snare operators users would describe themselves as professionals, there is no such thing as professional use. As an example, in a Defra-funded study into the use of snares, the Game & Wildlife Conservation Trust staff member who undertook the field trials *had 20 years experience as a field-based wildlife biologist, working almost entirely on projects requiring the detection, capture, handling, tracking or humane dispatch of mammalian predator species. He has considerable experience of using snares to catch foxes for radio-tagging studies and ... provides input into industry recognised best practice guidelines, training and other educational material on fox snaring* (Anon. 2012). Despite this expertise, 73% of his reported captures were non-target species (Harris 2022).

Elsewhere in the Defra-funded study, it is stated that a technician was used who was *fully competent in the use of snares because this focuses on the device as operated according to best practice (because it is difficult and unethical to emulate bad practice)*. In 211 snare nights in familiar areas, this technician caught 18 animals: 3 foxes, 13 hares, and 2 badgers. Five hares, 2 badgers and 1 fox escaped: 1 fox and 1 badger escaped with the snare attached. Three hares were severely injured/dead and 2 hares were predated. Both foxes that were held had haemorrhages on their necks extending into the deeper muscle. Although the snares were placed so as to avoid entanglement with fences or woody shrubs, three of the snares were entangled with non-woody vegetation (Anon. 2012). Assuming that the animals that escaped with the snare attached suffered significant adverse welfare impacts, 7/18 captures (39%) were severely injured, dead or predated; just 2 target species (11% of captures) were held and killed (Harris 2022).

There is no evidence that professional/highly experienced operators catch fewer non-target species than other users, or cause fewer deaths or injuries to both target and non-target species.

The Animal Welfare Act 2006 prohibits causing unnecessary suffering to a captive animal. There are very significant welfare concerns about the use of snares, even when set by professional/highly experienced operators.

Would it be ethical to use snares to catch foxes for research?

The Animals (Scientific Procedures) Act 1986 regulates the use of animals, both captive and free-living, in scientific research, and licenses are required for scientific purposes that may cause pain, suffering, distress or lasting harm to an animal. So a scientist is likely to require both a project and personal licence from the Home Office under the the Animals (Scientific Procedures) Act 1986 to undertake a study that involves catching, handling, sedating and/or marking a wild mammal (<http://www.homeoffice.gov.uk/scienceresearch/animal-research/>). Post 1986, all of my research that involved catching, handling, sedating and marking foxes was undertaken under Home Office licences. It also required approval from the University of Bristol's ethics committee – see below.

The Royal Society is the premier scientific organisation in Britain: its role is to recognise, promote, and support excellence in science. The Royal Society states that *all research should be carried out with a high regard for animal welfare*, and that *All research involving animals must have been reviewed and approved by an ethics committee prior to commencing the study and performed in accordance with relevant institutional and national guidelines and regulations* (<https://royalsociety.org/journals/ethics-policies/research-ethics/>).

The Royal Society of Edinburgh is Scotland's national academy: it states that good research conduct should only include *procedures that are aimed at avoiding unreasonable risk or harm to humans, animals or the environment* (<https://rse.org.uk/funding-collaboration/codes-of-conduct/>).

All reputable research institutions have an ethics committee: see, for instance those for the universities of Aberdeen, St Andrews and Stirling (https://www.University_of_Aberdeen_Statement_on_Use_of_Animals_in_Research.pdf, <https://www.ed.ac.uk/research/animal-research>, <https://www.st-andrews.ac.uk/research/integrity-ethics/animals/policy/> and <https://www.stir.ac.uk/research/research-ethics-and-integrity/animal-research-at-the-university-of-stirling/our-research-involving-animals/>). It is extremely unlikely that any ethics committee would approve a research project that involved the use of snares to catch foxes for research when the British Veterinary Association, the British Veterinary Zoological Society and Scottish Animal Welfare Commission have all concluded that the use of snares raises significant animal welfare concerns and called for an outright ban on their sale and use both by the general public and trained operators/industry.

Furthermore, there has been considerable pressure to reduce the number of animals used in scientific research. The underlying principle, referred to as the 3Rs (Replacement, Reduction and Refinement), provides a framework for performing more humane animal research. The 3Rs have been embedded in national and international legislation and regulations on the use of animals in scientific procedures, as well as in the policies of organisations that fund or conduct animal research (<https://www.nc3rs.org.uk/who-we-are/3rs>). See, for instance, the policies

of Aberdeen and Edinburgh universities

(<https://www.abdn.ac.uk/staffnet/documents/policy-zone-research-and-knowledge-exchange/RGH-Section-3.pdf> and <https://www.ed.ac.uk/research/animal-research/animal-welfare-ethics/ethical-review-and-awerb>).

It is extremely unlikely that the ethics committee of a reputable research organisation would sanction the use of snares to catch foxes for research.

Would studies that use snares to catch foxes meet ethical standards for publication?

As part of the drive to improve animal welfare standards in research, most reputable scientific journals have introduced ethics policies and/or ethical committees that review submitted papers and only consider them for publication if the research was conducted in accordance with the highest animal welfare standards.

For instance, the Association for the Study of Animal Behaviour (ASAB) and the Animal Behavior Society (ABS) have formed ethical and animal care committees, respectively, and each appoints an ethics editor to serve on the editorial board of their scientific journal *Animal Behaviour* (ASAB Ethical Committee/ABS Animal Care Committee 2023). Similarly, the *Journal of Mammalogy* states that investigators conducting research requiring live capture of mammals assume the responsibility for using humane methods that respect target and non-target species (Sikes and the Animal Care and Use Committee of the American Society of Mammalogists 2016).

It is unlikely that any reputable scientific journal would accept a paper for publication that used capture techniques such as snares that cause considerable harm to both target and non-target animals.

Conclusion

There is no scientific case to permit the use of snares to catch foxes for scientific research in Scotland.

References

Anon. (2012) *Determining the extent of use and humaneness of snares in England and Wales.*

http://sciencesearch.defra.gov.uk/Document.aspx?Document=9872_wm0315-humaneness-snares.pdf.

ASAB Ethical Committee/ABS Animal Care Committee (2023) Guidelines for the ethical treatment of nonhuman animals in behavioural research and teaching. *Animal Behaviour* **195**: i-xi.

British Veterinary Association (2022) Executive summary – BVA and BVZS position on the use and sale of snares for the control of free-ranging wildlife.

<https://www.bva.co.uk/media/4625/bva-and-bvzs-position-on-the-use-and-sale-of-snares-executive-summary.pdf>.

Dorning J & Harris S (2019) The challenges of recognising individuals with few distinguishing features: identifying red foxes *Vulpes vulpes* from camera-trap photos. *PloS One* **14**: e0216531.

Grove P & Oswald J (2001) Western capercaillie captures in snares. *Scottish Birds* **22**: 28-30.

Harris S (2022) *A review of the use of snares in the UK.*

<https://www.antisnaring.org.uk/news/nasc/uk-mammal-expert-ban-snaring-protect-animal-welfare/>.

Scottish Animal Welfare Commission (2022) *Trapping of terrestrial wild mammals using snares: position paper.* <https://www.gov.scot/publications/scottish-animal-welfare-commission-trapping-of-terrestrial-wild-mammals-using-snares-position-paper/>.

Sikes RS and the Animal Care and Use Committee of the American Society of Mammalogists (2016) 2016 guidelines of the American Society of Mammalogists for the use of wild mammals in research and education. *Journal of Mammalogy* **97**: 663-688.

White PCL, Newton-Cross GA, Moberly RL, Smart JC, Baker PJ & Harris S (2003) The current and future management of wild mammals hunted with dogs in England and Wales. *Journal of Environmental Management* **67**: 187-197.

Appendix: peer-reviewed scientific studies on foxes in Scotland published since 1970

1. Atterby H, Allnutt TR, MacNicoll AD, Jones EP & Smith GC (2015) Population genetic structure of the red fox (*Vulpes vulpes*) in the UK. *Mammal Research* **60**: 9-19.
2. Baines D, Moss R & Dugan D (2004) Capercaillie breeding success in relation to forest habitat and predator abundance. *Journal of Applied Ecology* **41**: 59-71.
3. Baines D, Warren P & Richardson M (2007) Variations in the vital rates of black grouse *Tetrao tetrix* in the United Kingdom. *Wildlife Biology* **13**: 109-116.
4. Baines D, Redpath S, Richardson M & Thirgood S (2008) The direct and indirect effects of predation by hen harriers *Circus cyaneus* on trends in breeding birds on a Scottish grouse moor. *Ibis* **150**: 27-36.
5. Baines D, Aebischer N, Macleod A & Woods J (2013) Pine marten *Martes martes* and red fox *Vulpes vulpes* sign indices in Scottish forests: population change and reliability of field identification of scats. *Wildlife Biology* **19**: 490-495.
6. Baines D, Aebischer NJ & Macleod A (2016) Increased mammalian predators and climate change predict declines in breeding success and density of Capercaillie *Tetrao urogallus*, an old stand specialist, in fragmented Scottish forests. *Biodiversity and Conservation* **25**: 2171-2186.
7. Baines D, Fletcher K, Hesford N, Newborn D & Richardson M (2023) Lethal predator control on UK moorland is associated with high breeding success of curlew, a globally near-threatened wader. *European Journal of Wildlife Research* **69(1)**: 6.
8. Bartley PM, Hamilton C, Wilson C, Innes EA & Katzer F (2016) Detection of *Babesia annae* DNA in lung exudate samples from red foxes (*Vulpes vulpes*) in Great Britain. *Parasites & Vectors* **9**: 1-8.

9. Beard PM, Henderson D, Daniels MJ, Pirie A, Buxton D, Greig A, Hutchings MR, McKendrick I, Rhind S, Stevenson K & Sharp JM (1999) Evidence of paratuberculosis in fox. *Veterinary Record* **145**: 612-613.
10. Beard PM, Daniels MJ, Henderson D, Pirie A, Rudge K, Buxton D, Rhind S, Greig A, Hutchings MR, McKendrick I, Stevenson K & Sharp JM (2001) Paratuberculosis infection of nonruminant wildlife in Scotland. *Journal of Clinical Microbiology* **39**: 1517-1521.
11. Calder, CJ & Gorman ML (1991) The effects of red fox *Vulpes vulpes* faecal odours on the feeding behaviour of Orkney voles *Microtus arvalis*. *Journal of Zoology* **224**: 599-606.
12. Daniels MJ, Hutchings MR, Beard PM, Henderson D, Greig A, Stevenson K & Sharp JM (2003). Do non-ruminant wildlife pose a risk of paratuberculosis to domestic livestock and vice versa in Scotland? *Journal of Wildlife Diseases* **39**: 10-15.
13. Davison A, Birks JDS, Brookes RC, Braithwaite TC & Messenger JE (2002) On the origin of faeces: morphological versus molecular methods for surveying rare carnivores from their scats. *Journal of Zoology* **257**: 141-143.
14. Dobson A & Hudson P (1995) The interaction between the parasites and predators of red grouse *Lagopus lagopus scoticus*. *Ibis* **137**: S87-S96.
15. Douglas DJ, Bellamy PE, Stephen LS, Pearce-Higgins JW, Wilson JD & Grant MC (2014) Upland land use predicts population decline in a globally near-threatened wader. *Journal of Applied Ecology* **51**: 194-203.
16. Douglas DJ, Tománková I, Gullett P, Dodd SG, Brown D, Clift M, Russell N, Warnock N, Smart J & Sanders S (2023) Varying response of breeding waders to experimental manipulation of their habitat and predators. *Journal for Nature Conservation* **72**: 126353.
17. Dyczkowski J & Yalden DW (1998) An estimate of the impact of predators on the British field vole *Microtus agrestis* population. *Mammal Review* **28**: 165-184.
18. Elliot RD (1985) The effects of predation risk and group size on the anti-predator responses of nesting lapwings *Vanellus vanellus*. *Behaviour* **92**: 168-187.
19. Evans DM, Villar N, Littlewood NA, Pakeman RJ, Evans SA, Dennis P, Skartveit J & Redpath SM (2015) The cascading impacts of livestock grazing in upland ecosystems: a 10-year experiment. *Ecosphere* **6**: 1-15.
20. Fox NJ, Smith LA, Stevenson K, Davidson RS, Marion G & Hutchings MR (2020) Infection of non-ruminant wildlife by *Mycobacterium avium* subsp. *paratuberculosis*. In Behr MA & Collins DM (eds) *Paratuberculosis: organism, disease, control*, pp. 200-212. CABI, Wallingford.
21. Frank LG (1979) Selective predation and seasonal variation in the diet of the fox (*Vulpes vulpes*) in NE Scotland. *Journal of Zoology* **189**: 526-532.
22. Giergiel M, Campbell S, Giela A, Sharp E, Casali F, Śniegocki T, Sell B & Jedziniak P (2023) Residues of an anthelmintic veterinary drug (closantel) detected in red foxes (*Vulpes vulpes*) in Scotland. *Ecotoxicology and Environmental Safety* **253**: 114651.
23. Green RE & Etheridge B (1999) Breeding success of the hen harrier *Circus cyaneus* in relation to the distribution of grouse moors and the red fox *Vulpes vulpes*. *Journal of Applied Ecology* **36**: 472-483.

24. Hamilton CM, Gray R, Wright SE, Gangadharan B, Laurenson K & Innes EA (2005) Prevalence of antibodies to *Toxoplasma gondii* and *Neospora caninum* in red foxes (*Vulpes vulpes*) from around the UK. *Veterinary Parasitology* **130**: 169-173.
25. Hamilton GA, Ruthven AD, Findlay E, Hunter K & Lindsay DA (1981) Wildlife deaths in Scotland resulting from misuse of agricultural chemicals. *Biological Conservation* **21**: 315-326.
26. Hancock MH, Klein D & Cowie NR (2020) Guild-level responses by mammalian predators to afforestation and subsequent restoration in a formerly treeless peatland landscape. *Restoration Ecology* **28**: 1113-1123.
27. Helm J, Roberts L, Jefferies R, Shaw SE & Morgan ER (2015) Epidemiological survey of *Angiostrongylus vasorum* in dogs and slugs around a new endemic focus in Scotland. *Veterinary Record* **177**: 46-51.
28. Hewson R (1976) A population study of mountain hares (*Lepus timidus*) in north-east Scotland from 1956-1969. *Journal of Animal Ecology* **45**: 395-414.
29. Hewson R (1981) Scavenging of mammal carcasses by birds in West Scotland. *Journal of Zoology* **194**: 525-537.
30. Hewson R (1983) The food of wild cats (*Felis silvestris*) and red foxes (*Vulpes vulpes*) in west and north-east Scotland. *Journal of Zoology* **200**: 283-289.
31. Hewson R (1984) Changes in the numbers of foxes (*Vulpes vulpes*) in Scotland. *Journal of Zoology* **203**: 561-569.
32. Hewson R (1984) Scavenging and predation upon sheep and lambs in west Scotland. *Journal of Applied Ecology* **21**: 843-868.
33. Hewson R (1985) Lamb carcasses and other food remains at fox dens in Scotland. *Journal of Zoology* **206**: 291-296.
34. Hewson R (1986) Distribution and density of fox breeding dens and the effects of management. *Journal of Applied Ecology* **23**: 531-538.
35. Hewson R (1994) The use of dens by hill foxes (*Vulpes vulpes*). *Journal of Zoology* **233**: 331-335.
36. Hewson R (1995) Use of salmonid carcasses by vertebrate scavengers. *Journal of Zoology* **235**: 53-65.
37. Hewson R & Kolb HH (1973) Changes in the numbers and distribution of foxes (*Vulpes vulpes*) killed in Scotland from 1948-1970. *Journal of Zoology* **171**: 345-365.
38. Hewson R & Kolb HH (1974) The control of foxes in Scottish forests. *Scottish Forestry* **28**: 272-282.
39. Hewson R & Kolb HH (1975) The food of foxes (*Vulpes vulpes*) in Scottish forests. *Journal of Zoology* **176**: 287-292.
40. Hewson R & Kolb HH (1976) Scavenging on sheep carcasses by foxes (*Vulpes vulpes*) and badgers (*Meles meles*). *Journal of Zoology* **180**: 496-498.
41. Hewson R & Leitch AF (1983) The food of foxes in forests and on the open hill. *Scottish Forestry* **37**: 39-50.
42. Hewson R & Verkaik AJ (1981) Body condition and ranging behaviour of blackface hill sheep in relation to lamb survival. *Journal of Applied Ecology* **18**: 401-415.

43. Hewson R & Wilson CJ (1979) Home range and movements of Scottish Blackface sheep in Lochaber, north-west Scotland. *Journal of Applied Ecology* **16**: 743-751.
44. Hudson PJ & Dobson AP (1995) Extrinsic factors influencing the population dynamics of red grouse, *Lagopus lagopus scoticus*. In Jenkins D (ed.) *Proceedings of the 6th International Grouse Symposium*, pp. 19-26. World Pheasant Association, Reading.
45. Hudson PJ, Newborn D & Robertson PA (1997) Geographical and seasonal patterns of mortality in red grouse *Lagopus lagopus scoticus* populations. *Wildlife Biology* **3**: 79-87.
46. Jahren T, Storaas T, Willebrand T, Moa P F & Hagen BR (2016) Declining reproductive output in capercaillie and black grouse - 16 countries and 80 years. *Animal Biology* **66**: 363-400.
47. Kirk L, Limon G, Guitian FJ, Hermosilla C & Fox MT (2014) *Angiostrongylus vasorum* in Great Britain: a nationwide postal questionnaire survey of veterinary practices. *Veterinary Record* **175**: 118-123.
48. Kolb H (1978) The formation of lines in the cementum of premolar teeth in foxes. *Journal of Zoology* **185**: 259-263.
49. Kolb HH (1978) Variation in the size of foxes in Scotland. *Biological Journal of the Linnean Society* **10**: 291-304.
50. Kolb HH (1984) Factors affecting the movements of dog foxes in Edinburgh. *Journal of Applied Ecology* **21**: 161-173.
51. Kolb HH (1985) Habitat use by foxes in Edinburgh. *Revue d'Écologie (La Terre et La Vie)* **40**: 139-143.
52. Kolb HH (1986) Some observations on the home ranges of vixens (*Vulpes vulpes*) in the suburbs of Edinburgh. *Journal of Zoology* **210**: 636-639.
53. Kolb HH & Hewson R (1974) The body size of the red fox (*Vulpes vulpes*) in Scotland. *Journal of Zoology* **173**: 253-255.
54. Kolb HH & Hewson R (1979) Variation in the diet of foxes in Scotland. *Acta Theriologica* **24**: 69-83.
55. Kolb HH & Hewson R (1980) A study of fox populations in Scotland from 1971 to 1976. *Journal of Applied Ecology* **17**: 7-19.
56. Kolb HH & Hewson R (1980) The diet and growth of fox cubs in two regions of Scotland. *Acta Theriologica* **25**: 325-331.
57. Lambin X, Petty SJ & Mackinnon JL (2000) Cyclic dynamics in field vole populations and generalist predation. *Journal of Animal Ecology* **69**: 106-119.
58. Learmount J, Zimmer IA, Conyers C, Boughtflower VD, Morgan CP & Smith GC (2012) A diagnostic study of *Echinococcus multilocularis* in red foxes (*Vulpes vulpes*) from Great Britain. *Veterinary Parasitology* **190**: 447-453.
59. Leckie FM, Thirgood SJ, May R & Redpath SM (1998) Variation in the diet of red foxes on Scottish moorland in relation to prey abundance. *Ecography* **21**: 599-604.
60. Littlewood NA, Mason TH, Hughes M, Jaques R, Whittingham MJ & Willis SG (2019) The influence of different aspects of grouse moorland management on nontarget bird assemblages. *Ecology and Evolution* **9**: 11,089-11,101.
61. Lockie J (1973) Wildlife and land use in highland Scotland. *Oryx* **12**: 53-63.

62. Ludwig SC, Roos S, Bubb D & Baines D (2017) Long-term trends in abundance and breeding success of red grouse and hen harriers in relation to changing management of a Scottish grouse moor. *Wildlife Biology* **2017**: wlb.00246.
63. Ludwig SC, Roos S & Baines D (2019) Responses of breeding waders to restoration of grouse management on a moor in south-west Scotland. *Journal of Ornithology* **160**: 789-797.
64. Macdonald DW, Bunce RGH & Bacon PJ (1981) Fox populations, habitat characterization and rabies control. *Journal of Biogeography* **8**: 145-151.
65. McMillan RL (2014) Hen harriers on Skye, 2000-12: nest failures and predation. *Scottish Birds* **34**: 30-39.
66. Meredith AL, Cleaveland SC, Brown J, Mahajan A & Shaw DJ (2015) Seroprevalence of *Encephalitozoon cuniculi* in wild rodents, foxes and domestic cats in three sites in the United Kingdom. *Transboundary and Emerging Diseases* **62**: 148-156.
67. Moberly RL, White PCL, Webbon CC, Baker PJ & Harris S (2003) Factors associated with fox (*Vulpes vulpes*) predation of lambs in Britain. *Wildlife Research* **30**: 219-227.
68. Moberly RL, White PC, Webbon CC, Baker PJ & Harris S (2004) Modelling the costs of fox predation and preventive measures on sheep farms in Britain. *Journal of Environmental Management* **70**: 129-143.
69. Morgan ER, Tomlinson A, Hunter S, Nichols T, Roberts E, Fox MT & Taylor MA (2008) *Angiostrongylus vasorum* and *Eucoleus aerophilus* in foxes (*Vulpes vulpes*) in Great Britain. *Veterinary Parasitology* **154**: 48-57.
70. Moss R, Trenholm I B, Watson A & Parr R (1990) Parasitism, predation and survival of hen red grouse *Lagopus lagopus scoticus* in spring. *Journal of Animal Ecology* **59**: 631-642.
71. Naylor JR & Knott JG (2018) A pack of dogs is more effective at flushing red foxes to guns than a pair. *Wildlife Society Bulletin* **42**: 338-346.
72. Newey S, Dahl F, Willebrand T & Thirgood S (2007) Unstable dynamics and population limitation in mountain hares. *Biological Reviews* **82**: 527-549.
73. O'Mahony D, Lambin X, MacKinnon JL & Coles CF (1999) Fox predation on cyclic field vole populations in Britain. *Ecography* **22**: 575-581.
74. Parr R (1992) The decline to extinction of a population of golden plover in north-east Scotland. *Ornis Scandinavica* **23**: 152-158.
75. Parr R (1993) Nest predation and numbers of golden plovers *Pluvialis apricaria* and other moorland waders. *Bird Study* **40**: 223-231.
76. Patterson IJ (1977) The control of fox movement by electric fencing. *Biological Conservation* **11**: 267-278.
77. Pettett CE, Johnson PJ, Moorhouse TP & Macdonald DW (2018) National predictors of hedgehog *Erinaceus europaeus* distribution and decline in Britain. *Mammal Review* **48**: 1-6.
78. Philbey AW & Delgado D (2013) Detection of *Angiostrongylus vasorum* in red foxes in Scotland. *Veterinary Record* **173(6)**: 148.
79. Ratcliffe PR (1970) The occurrence of vestigial teeth in badger (*Meles meles*), roe deer (*Capreolus capreolus*) and fox (*Vulpes vulpes*) from the county of Argyll, Scotland. *Journal of Zoology* **162**: 521-525.

80. Scott DM, Berg MJ, Tolhurst BA, Chauvenet ALM, Smith GC, Neaves K, Lochhead J & Baker PJ (2014) Changes in the distribution of red foxes (*Vulpes vulpes*) in urban areas in Great Britain: findings and limitations of a media-driven nationwide survey. *PLoS One* **9(6)**: e99059.
81. Scott DM, Baker R, Tomlinson A, Berg MJ, Charman N & Tolhurst BA (2020) Spatial distribution of sarcoptic mange (*Sarcoptes scabiei*) in urban foxes (*Vulpes vulpes*) in Great Britain as determined by citizen science. *Urban Ecosystems* **23**: 1127-1140.
82. Smith GC, Gangadharan B, Taylor Z, Laurenson MK, Bradshaw H, Hide G, Hughes JM, Dinkel A, Romig T & Craig PS (2003) Prevalence of zoonotic important parasites in the red fox (*Vulpes vulpes*) in Great Britain. *Veterinary Parasitology* **118**: 133-142.
83. Soulsbury CD, Iossa G, Baker PJ, Cole NC, Funk SM & Harris S (2007) The impact of sarcoptic mange *Sarcoptes scabiei* on the British fox *Vulpes vulpes* population. *Mammal Review* **37**: 278-296.
84. Summers RW, Green RE, Proctor R, Dugan D, Lambie D, Moncrieff R, Moss R & Baines D (2004) An experimental study of the effects of predation on the breeding productivity of capercaillie and black grouse. *Journal of Applied Ecology* **41**: 513-525.
85. Summers RW, Willi J & Selvidge J (2009) Capercaillie *Tetrao urogallus* nest loss and attendance at Abernethy Forest, Scotland. *Wildlife Biology* **15**: 319-327.
86. Sumption KJ & Flowerdew JR (1985) The ecological effects of the decline in rabbits (*Oryctolagus cuniculus* L.) due to myxomatosis. *Mammal Review* **15**: 151-186.
87. Taylor CS, Gato RG, Learmount J, Aziz NA, Montgomery C, Rose H, Coulthwaite CL, McGarry JW, Forman DW, Allen S, Wall R & Morgan ER (2015) Increased prevalence and geographic spread of the cardiopulmonary nematode *Angiostrongylus vasorum* in fox populations in Great Britain. *Parasitology* **142**: 1190-1195.
88. Taylor MJ, Nevison I, Casali F, Giergiel M, Giela A, Campbell S, Sharp EA, Harley G & Posyniak A (2020) Investigation of the distribution of anticoagulant rodenticide residues in red fox (*Vulpes vulpes*) livers to ensure optimum sampling protocol. *Environmental Chemistry and Ecotoxicology* **2**: 50-55.
89. Thompson H, O'Keeffe AM, Lewis JCM, Stocker LR, Laurenson MK & Philbey AW (2010) Infectious canine hepatitis in red foxes (*Vulpes vulpes*) in the United Kingdom. *Veterinary Record* **166**: 111-114.
90. Thompson RCA (1976) The occurrence of *Mesocestoides* sp. in British wild red foxes (*Vulpes vulpes crucigera*). *Journal of Helminthology* **50**: 91-94.
91. Villar N, Lambin X, Evans D, Pakeman R & Redpath S (2013) Experimental evidence that livestock grazing intensity affects the activity of a generalist predator. *Acta Oecologica* **49**: 12-16.
92. Waggershauser CN, Taberlet P, Coissac E, Kortland K, Hambly C & Lambin X (2022) Interspecific coprophagia by wild red foxes: DNA metabarcoding reveals a potentially widespread form of commensalism among animals. *Ecology and Evolution* **12**: e9029.

93. Walker D, Abbondati E, Cox AL, Mitchell GBB, Pizzi R, Sharp CP & Philbey AW (2016) Infectious canine hepatitis in red foxes (*Vulpes vulpes*) in wildlife rescue centres in the UK. *Veterinary Record* **178**: 421-421.
94. Walker D, Fee SA, Hartley G, Learmount J, O'Hagan MJ, Meredith AL, Bronsvort BM de C, Porphyre T, Sharp CP & Philbey AW (2016) Serological and molecular epidemiology of canine adenovirus type 1 in red foxes (*Vulpes vulpes*) in the United Kingdom. *Scientific Reports* **6(1)**: 36051.
95. Walton KC (1984) Fluoride in fox bone near an aluminium reduction plant in Anglesey, Wales, and elsewhere in the United Kingdom. *Environmental Pollution Series B* **7**: 273-280.
96. Webbon CC, Baker PJ & Harris S (2004) Faecal density counts for monitoring changes in red fox numbers in rural Britain. *Journal of Applied Ecology* **41**: 768-779.
97. White PCL, Groves HL, Savery JR, Conington J & Hutchings MR (2000) Fox predation as a cause of lamb mortality on hill farms. *Veterinary Record* **147**: 33-37.
98. Wilson WL (1990) Changes in the diet of foxes (*Vulpes vulpes*) on the sands of Forvie National Nature Reserve, during the eider nesting period, 1974 to 1988. *Journal of Zoology* **221**: 305-308.
99. Wolfe A & Long AM (1997) Distinguishing between the hair fibres of the rabbit and the mountain hare in scats of the red fox. *Journal of Zoology* **242**: 370-375.
100. Zalewska K, Waggershauser C N, Kortland K & Lambin X (2021). The best defence is not being there: avoidance of larger carnivores is not driven by risk intensity. *Journal of Zoology* **315**: 110-122.